

# MONOPOLES

MICHAEL K. MURRAY

## 1. INTRODUCTION

Monopoles are solutions of a first order partial differential equation — the Bogomolny equation. They can be thought of as approximated by static, magnetic particles in  $\mathbb{R}^3$ . In these notes we will consider what monopoles are and some of the various approaches to understanding them. In particular we will discuss the spectral curve, twistor theory, Nahm transform and rational map of a monopole. A good starting reference for all this material is the book by Atiyah and Hitchin [2] on monopole scattering, in particular Chapter 16 ‘Background Material’ from which these notes have borrowed notation. See also the excellent survey by Sutcliffe [22].

## 2. MONOPOLES IN $\mathbb{R}^3$ .

We start with a Lie algebra which we will assume is  $\mathfrak{su}(2)$ , the Lie algebra of all skew-hermitian 2 by 2 matrices. Let  $A$  be a one-form with values in  $\mathfrak{su}(2)$  so that  $A = \sum_{i=1}^3 A_i dx^i$  and each  $A_i$  is a function  $A_i: \mathbb{R}^3 \rightarrow \mathfrak{su}(2)$ . The Higgs field  $\Phi$  is a function  $\Phi: \mathbb{R}^3 \rightarrow \mathfrak{su}(2)$ . The one-form  $A$  can be thought of as the connection one-form for a connection  $\nabla = d + A$  on a trivial  $SU(2)$  bundle on  $\mathbb{R}^3$ . The curvature of such a connection is the two-form

$$F_A = \frac{1}{2} \sum_{i,j=1}^3 F_{ij} dx^i \wedge dx^j$$

where

$$F_{ij} = [\nabla_i, \nabla_j] = \partial_i A_j - \partial_j A_i + [A_i, A_j].$$

The connection  $A$  can be used to covariantly differentiate the Higgs field  $\Phi$  to obtain

$$\nabla_A \Phi = \sum_{i=1}^3 (\partial_i \Phi + [A_i, \Phi]) dx^i.$$

A monopole is a connection  $A$  and a Higgs field  $\Phi$  satisfying the Bogomolny equations and some particular boundary conditions. The Bogomolny equations are:

$$(1) \quad F_A = * \nabla_A \Phi$$

where  $*$  is the Hodge star or duality operator from one-forms to two-forms defined by

$$\begin{aligned} *dx^1 &= dx^2 \wedge dx^3 \\ *dx^2 &= dx^3 \wedge dx^1 \\ *dx^3 &= dx^1 \wedge dx^2. \end{aligned}$$

If  $A$  and  $B$  are elements of  $\mathfrak{su}(2)$  let  $\langle A, B \rangle$  denote the invariant form  $\langle A, B \rangle = -\text{tr}(AB^t)$ . Then the energy density of a pair  $(A, \Phi)$  is defined by

$$(2) \quad e(A, \Phi) = \frac{1}{2}|F_A|^2 + \frac{1}{2}|\nabla_A \Phi|^2$$

where

$$|F_A|^2 = \sum_{i < j} \langle F_{ij}, F_{ij} \rangle$$

and

$$|\nabla_A \Phi|^2 = \frac{1}{2} \sum_i \langle \nabla_i \Phi, \nabla_i \Phi \rangle.$$

The Yang-Mills-Higgs action of a pair  $(A, \Phi)$  is the integral over three space of the energy density:

$$(3) \quad \mathcal{L}(A, \Phi) = \int_{\mathbb{R}^3} e(A, \Phi) d^3x.$$

If  $B_R$  is a ball of radius  $R$  integrating by parts shows that

$$\int_{B_R} e(A, \Phi) d^3x = \int_{B_R} \frac{1}{2} |F_A \pm * \nabla_A \Phi|^2 d^3x \mp \int_{S_R^2} \langle F_A, \Phi \rangle$$

where  $S_R^2$  is the sphere of radius  $R$ . If the limits of all these integrals exist as  $R \rightarrow \infty$  we obtain

$$\mathcal{L}(A, \Phi) = \int_{\mathbb{R}^3} \frac{1}{2} |F_A \pm * \nabla_A \Phi|^2 d^3x \mp \lim_{R \rightarrow \infty} \int_{S_R^2} \langle F_A, \Phi \rangle.$$

From this we easily deduce that the minima of the Yang-Mills-Higgs functional are solutions of the Bogomolny equations (1) or the anti-Bogomolny equations  $F_A = -* \nabla_A \Phi$ . As changing  $\Phi$  to  $-\Phi$  changes a solution of the Bogomolny equations to a solution of the anti-Bogomolny equations we concentrate our attention on the former.

The Bogomolny equations are invariant under *gauge transformations* that is replacing  $(A, \Phi)$  by

$$(gAg^{-1} + gd(g^{-1}), g\Phi g^{-1})$$

where  $g: \mathbb{R}^3 \rightarrow SU(2)$ . The energy density (2) is also invariant under gauge transformations. When we talk about a monopole we are really talking about an equivalence class of  $(A, \Phi)$  under gauge transformations.

The boundary conditions imposed on a monopole are primarily that the energy density (2) should have finite integral — that is the action (3) is finite. There are some additional technical conditions that we will not be concerned with. It is believed, in any case, that these can all be deduced from finiteness of the action and the Bogomolny equations. From these boundary conditions we can deduce that, after a suitable gauge transformation, we can arrange for the Higgs field to have a limiting value at infinity

$$\Phi^\infty(u) = \lim_{t \rightarrow \infty} \Phi(tu)$$

where  $u \in S^2$ . The boundary conditions can be used to show that the eigenvalues of the Higgs field at infinity are independent of the direction  $u \in S^2$ . In the case of  $SU(2)$  this is equivalent to the fact that  $|\Phi(u)|^2 = \text{constant}$  for all  $u$ .

It is easy to show that if  $c > 0$  and  $(A, \Phi)$  solves the Bogomolny equations then

$$(\hat{A}(x), \hat{\Phi}(x)) = (cA(x/c), c\Phi(x/c))$$

also solves the Bogomolny equations. So we may as well normalise the Higgs field so that  $|\Phi(u)|^2 = 1$  for all directions  $u$ . Because the Lie algebra  $\mathfrak{su}(2)$  is three dimensional the Higgs field at infinity is a map

$$\Phi^\infty: S^2 \rightarrow S^2 \subset \mathfrak{su}(2).$$

The space of all continuous maps  $S^2 \rightarrow S^2$  breaks up into connected components labelled by a winding number  $k \in \mathbb{Z}$  just as for maps  $S^1 \rightarrow S^1$ .

Because of this boundary condition we can arrange by a gauge transformation for any Higgs field to satisfy

$$\lim_{t \rightarrow \infty} \Phi(0, 0, t) = \frac{1}{\sqrt{2}} \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}.$$

We call such a Higgs field *framed*. We define the moduli space of all monopoles of charge  $k$ ,  $\mathcal{M}_k$ , to be the space of all  $(A, \Phi)$  solving the Bogomolny equations, satisfying the appropriate boundary conditions, with the Higgs field framed and modulo the action of gauge transformations satisfying

$$(4) \quad \lim_{t \rightarrow \infty} g(0, 0, t) = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

If  $k \leq 0$  then  $\mathcal{M}_k = \emptyset$ , otherwise  $\mathcal{M}_k$  is a manifold of dimension  $4k$ .

Notice that the Bogomolny equations are translation invariant. Moreover, because of the way we have defined the framing, the group of all diagonal matrices (a copy of the circle group  $S^1$ ) acts by constant gauge transformations on  $\mathcal{M}_k$ . Hence  $\mathbb{R}^3 \times S^1$  acts on  $\mathcal{M}_k$ .

If  $k = 1$  there is, up to this action of  $\mathbb{R}^3 \times S^1$ , a unique monopole, the Bogomolny, Prasad, Sommerfield (BPS) monopole, given by

$$\begin{aligned} \Phi(x) &= \left( \frac{1}{r} - \frac{1}{\tanh r} \right) \frac{e}{r} \\ A(x) &= \left( \frac{1}{\sinh r} - \frac{1}{r} \right) \frac{[e, de]}{r} \end{aligned}$$

where  $r = |x|$  and  $e(x) = \sum_{i=1}^3 x^i e^i$  for an orthonormal basis  $e^1, e^2, e^3$  of  $\mathfrak{su}(2)$ .

The other  $k = 1$  monopoles are obtained by acting by  $\mathbb{R}^3 \times S^1$  so that  $\mathcal{M}_1 = \mathbb{R}^3 \times S^1$ . It is easy to calculate the energy density of the BPS monopole as follows. For any monopole there is a useful formula

$$e(A, \Phi) = \sum_{i=1}^3 \frac{\partial^2}{(\partial x^i)^2} \langle \Phi, \Phi \rangle$$

which can be proved using the Bogomolny equations and the Bianchi identity. From this we find that for the BPS monopole:

$$e(A, \Phi) = \frac{6}{\tanh^4 r} - \frac{8}{\tanh^2 r} + 2 + \frac{2}{r^4} - \frac{8}{r \tanh^3 r} + \frac{8}{r \tanh r}$$

where  $r = |x|$ . Clearly the energy density is spherically symmetric about the origin in  $\mathbb{R}^3$ . If we plot it along the  $x$ -axis we obtain Figure 2 and we see that the energy is concentrated around the origin. We think of the monopole as a particle located at the origin.

The BPS monopole was discovered by Prasad and Sommerfield in 1975 [21]. For some time this was the only monopole known and it was unclear whether higher charge monopoles existed. In 1977 Manton [15] showed that to first order the forces




FIGURE 1. Energy density for  $k = 1$ , BPS monopole.

between two monopoles, due the Higgs field and the connection, cancelled. This lead to the conjecture that stable higher charge monopoles would exist. Weinberg in 1979 [23] calculated that the dimension of the moduli space of monopoles would be  $4k$  if it was non-empty. Finally in 1981 Taubes [12] showed that the moduli space was non-empty.

It is useful to briefly discuss Taubes' proof as it reveals some of the interesting structure of the moduli space. Taubes started with  $k$  points a long way apart in  $\mathbb{R}^3$  surrounded by very large balls. At the centre of each of these balls he placed a BPS monopole and joined these all together smoothly. The result is not, of course, a solution to the Bogomolny equations but it is close to one. Such an approximate solution can be used as the initial stage of an iterative scheme which converges to a real solution.

The solutions produced by Taubes are on the 'edge' of the moduli space. That is, the moduli space  $\mathcal{M}_k$  has a compact region outside of which the monopoles approximate a superposition BPS monopoles at  $k$  points. In particular the energy density will be concentrated at these  $k$  points. One might also expect that the Higgs field has a zero at each of these  $k$  points but this has not been proved. Note that thinking of the charge  $k$  monopoles as  $k$  BPS monopoles gives an explanation of the dimension of the moduli space: each constituent BPS monopole contributes its position (three parameters) and a phase, which is a point on the circle, making a total of four parameters.

It is important to note that when the charge is greater than one we cannot associate to every monopole  $(A, \Phi)$  a collection of  $k$  points which are the locations of the  $k$  particles we think of as its constituents. We can however associate sensibly to a  $k$  monopole a *centre of mass* or location [8] which we define below.

The analysis of monopoles directly in terms of the connection and Higgs field on  $\mathbb{R}^3$ , for example the definition of the location of a monopole, while possible, is difficult. Part of this difficulty stems from the infinite dimensional symmetry group of gauge transformations. Research on monopoles has focused on various transformations which are designed to construct some other mathematical data equivalent to the monopole. Study of these data then, hopefully, sheds light on the original monopole. This process is particularly useful if the object produced is an invariant of the monopole, something which does not change under gauge transformation. The transforms we will discuss below are:

- the spectral curve which is an invariant of the monopole,
- Hitchin's adaption of the twistor transform which relates monopoles to holomorphic bundles on mini-twistor space,
- the Nahm transform which relates monopoles to solutions of an ordinary differential equation on the interval  $(-1, 1)$ , and
- the rational map which is another invariant of the monopole.

### 3. THE SPECTRAL CURVE

Let  $\gamma(t)$  be an oriented line in  $\mathbb{R}^3$ . This can be put in the form

$$\gamma(t) = v + tu$$

with vectors  $u$  and  $v$  determined uniquely by the requirement that  $|u| = 1$ ,  $\langle u, v \rangle = 0$  and  $u$  points in the direction of the orientation. Along the line  $\gamma$  we have Hitchin's differential equation [6]

$$(5) \quad \left( \frac{d}{dt} + \sum_{i=1}^3 u^i A_i(\gamma(t)) - i\Phi(\gamma(t)) \right) s(t) = 0.$$

This is an ordinary differential equation so it has a two-dimensional space of solutions  $E_\gamma$  depending, of course, on  $\gamma$ .

Notice that by a gauge transformation we can arrange, for any given line  $\gamma$ , that  $\sum_{i=1}^3 u^i A_i = 0$  so we can essentially disregard this term. The boundary conditions can be used to show that we can expand the Higgs field along any line as

$$\Phi(\gamma(t)) = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix} - \frac{1}{2t} \begin{pmatrix} ik & 0 \\ 0 & -ik \end{pmatrix} + O\left(\frac{1}{t^2}\right)$$

where the  $k$  here is the monopole charge. We want to consider the Hitchin equation as a perturbation of a modified Hitchin equation:

$$(6) \quad \left( \frac{d}{dt} + \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} - \frac{1}{2t} \begin{pmatrix} k & 0 \\ 0 & -k \end{pmatrix} \right) s = 0.$$

The modified equation is the Hitchin equation with the  $o(1/t^2)$  term in the Higgs field expansion set to zero. We use this to study the behaviour of solutions of the Hitchin equation. The solutions of (6) are given by

$$(7) \quad s_1(t) = \begin{pmatrix} t^{k/2} e^{-t} \\ 0 \end{pmatrix}$$

$$(8) \quad s_2(t) = \begin{pmatrix} 0 \\ t^{-k/2} e^t \end{pmatrix}.$$

Asymptotic analysis of ordinary differential equations shows that for any line there are solutions  $s_1(t)$  and  $s_2(t)$  of the Hitchin equation (5) which behave asymptotically like the solutions to the modified Hitchin equation, that is they satisfy

$$\lim_{t \rightarrow \infty} t^{-k/2} e^t s_1(t) = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$\lim_{t \rightarrow \infty} t^{k/2} e^{-t} s_2(t) = \begin{pmatrix} 0 \\ 1 \end{pmatrix}.$$

Similarly there are solutions that decay and blow up exponentially as  $t \rightarrow -\infty$ . For the modified Hitchin equation a solution that blows up (decays) at one end of the line decays (blows up) at the other end. In general this will not be true of solutions of the Hitchin equation. In particular asymptotic analysis tells us that there will be a ball in  $\mathbb{R}^3$  of radius  $R > 0$  with the property that if a line lies outside the ball then the solutions  $s_1(t)$  and  $s_2(t)$  behave like the solutions to the modified Hitchin equation, that is  $s_1(t)$  decays as  $t \rightarrow -\infty$  and  $s_2(t)$  blows up as  $t \rightarrow -\infty$ .

We expect then that lines which *do not* exhibit this behaviour are somehow close to the monopole. We call a line  $\gamma$  a *spectral line* if there is a solution to the Hitchin equation which decays at both ends. We call the set of all spectral lines the *spectral curve* of the monopole. It is easy to see that being a spectral line for a monopole is independent of gauge transformations so the spectral curve is an invariant of the monopole.

It is not difficult to show that for the BPS monopole located at the point  $x \in \mathbb{R}^3$  the spectral lines are exactly the lines passing through  $x$ . Note that this is a two-dimensional set, indeed a copy of  $S^2$ . This is more generally true: the spectral curve is always a two-dimensional family of lines. To say more about the structure of the spectral curve we need to consider the set of all oriented lines in  $\mathbb{R}^3$ .

The importance of the spectral curve is the following theorem of Hitchin:

**Theorem 3.1** (Hitchin [6]). *If monopoles  $(A, \Phi)$  and  $(A', \Phi')$  have spectral curves  $S$  and  $S'$  and  $S = S'$  then  $(A, \Phi)$  is an unframed gauge transform of  $(A', \Phi')$ .*

Note that  $S = S'$  means equality of the corresponding sets of oriented lines in  $\mathbb{R}^3$  and that an unframed gauge transformation is one without the requirement that  $\lim_{t \rightarrow \infty} g(0, 0, t) = 1$  (c.f. equation (4)). Theorem 3.1 tells us that the spectral curve determines the monopole up to the action of the circle on the moduli space. It is possible to also add a small amount of additional data to the spectral curve which captures this phase, we shall not concern ourselves with that here.

#### 4. THE TWISTOR THEORY OF MONOPOLES

As we have discussed above each oriented line in  $\mathbb{R}^3$  is determined uniquely by vectors  $u$  and  $v$  satisfying  $|u| = 1$ ,  $\langle u, v \rangle = 0$ . It follows that the set of all oriented lines is the tangent bundle to the two-sphere:

$$TS^2 = \{(u, v) \mid |u| = 1, \langle u, v \rangle = 0\}.$$

This is often called the *mini-twistor* space of  $\mathbb{R}^3$ . Mini-twistor space is naturally a complex manifold and we can introduce co-ordinates  $(\eta, \zeta)$  on the open subset where  $u \neq (0, 0, 1)$  by letting

$$(9) \quad \zeta = \frac{u^1 + iu^2}{1 - u^3} \quad \text{and} \quad \eta = (v^1 + iv^2) + 2v^3\zeta + (-v^1 + iv^2)\zeta^2.$$

The relationship between mini-twistor space and  $\mathbb{R}^3$  is summarised by the equation:

$$(10) \quad \eta = (x^1 + ix^2) + 2x^3\zeta + (-x^1 + ix^2)\zeta^2.$$

If we hold  $(\eta, \zeta)$  fixed then the  $(x^1, x^2, x^3)$  satisfying (10) define a line in  $\mathbb{R}^3$ . On the other hand if we hold  $(x^1, x^2, x^3)$  fixed then the  $(\eta, \zeta)$  satisfying (10) parametrise the set of all lines through the point  $x = (x^1, x^2, x^3)$  which is a copy of  $S^2$  inside  $TS^2$ .

Mini-twistor space has an involution  $\tau: TS^2 \rightarrow TS^2$  which sends each oriented line to the same line with opposite orientation. In co-ordinates this is given by:

$$\tau(\eta, \zeta) = \left( -\frac{\bar{\eta}}{\bar{\zeta}^2}, -\frac{1}{\bar{\zeta}} \right).$$

Because  $\tau$  is like a conjugation it is called the *real structure*. The set of all lines through the point  $x$  is real in the sense that it is fixed by the real structure.

We can now state the basic result concerning the spectral curve. It is a subset of  $TS^2$  defined by an equation of the form

$$(11) \quad p(\eta, \zeta) = \eta^k + a_1(\zeta)\eta^{k-1} + \dots + a_k(\zeta) = 0$$

where each of the  $a_i(\zeta)$  is a polynomial of degree  $2i$ .

Note that by no means every such curve is the spectral curve of a monopole. One constraint is immediate from our definition. If a line is a spectral line then so also is the line with the opposite orientation. So the spectral curve is *real*, that is, fixed by the real structure. But more is true. The family of real curves defined by equations of the form (11) is  $(k-1)^2 - 1$  real dimensional whereas the moduli space of monopoles is  $4k$  dimensional. So there have to be further constraints on the spectral curve. In particular a certain holomorphic line bundle must be trivial when restricted to the spectral curve. It is possible to say quite precisely what the other constraints are and hence to say, in principle, which spectral curves give rise to monopoles [7].

Spectral curves can be used to deduce a number of useful facts about monopoles. It is easy to show that the only real curves of the type (11) for  $k=1$  are those of the form (10) for some point  $(x^1, x^2, x^3)$ . Hence the BPS monopoles are the only charge one monopoles.

The coefficient  $a_1(\zeta)$  in (11) defines a real curve and hence has the form

$$a_1(\zeta) = (x^1 + ix^2) + 2x^3\zeta + (-x^1 + ix^2)\zeta^2$$

for some point  $(x^1, x^2, x^3)$  in  $\mathbb{R}^3$ . This point is called the centre of the monopole. If  $(u, v_1), \dots, (u, v_k)$  is a collection of  $k$  parallel lines let us define their average to be the line  $(u, (1/k) \sum_{i=1}^k v_i)$ . Notice from definition of  $\eta$  (9) that if these lines have complex co-ordinates  $(\eta_1, \zeta), \dots, (\eta_k, \zeta)$  then their average has complex co-ordinates  $((1/k) \sum_{i=1}^k \eta_i, \zeta)$ . If we fix a particular direction in  $\mathbb{R}^3$ , that is fix a  $\zeta$  and look for all the spectral lines in that direction we are finding all the  $\eta$  satisfying a degree  $k$  polynomial and hence there are generically  $k$  of them. If we take the average of all these lines then it will pass through the monopole centre. This gives us a way of defining the centre entirely in  $\mathbb{R}^3$ . Take the average of the spectral lines in each direction in  $\mathbb{R}^3$ , the resulting family of lines will (nearly) all intersect in a single point, that point is the centre of the monopole.

The definition of the spectral curve of a monopole clearly preserves the action of the rotations and translations of  $\mathbb{R}^3$  and this gives us a way of looking for monopoles

with particular symmetries. We look first for spectral curves with these symmetries. This approach can be used to show that the only spherically symmetric monopole is the BPS monopole at the origin. It was used by Hitchin [7] to classify the axially symmetric monopoles and more recently [8] to find monopoles with symmetry groups those of the regular solids.

The various properties of the spectral curve such as the form of equation (11) are proved by using Hitchin's twistor transform. Hitchin [6] introduces the vector space  $E_\gamma$  of all solutions to his equation (5). This is a two-dimensional space and the collection of them all defines a complex vector bundle  $E$  over the mini-twistor space  $TS^2$ . Hitchin shows that the Bogomolny equations imply that  $E$  is a holomorphic vector bundle and moreover the monopole can be recovered from knowing  $E$ . The boundary conditions of the monopole then enter by noting that there are two distinguished holomorphic sub-bundles  $E^+$  and  $E^-$  of solutions to Hitchin's equation which decay at  $+$  and  $-$  infinity. The spectral curve is the set where these line bundles coincide. Algebraic geometry can then be used to prove Theorem 3.1 and that the spectral curve satisfies an equation of the form (11). Various constraints on the spectral curve also follow from the twistor theory. The precise constraints that a curve must satisfy to be the spectral curve are given in [7]. The proof that a spectral curve satisfying these constraints comes from a monopole requires the Nahm transform which we consider next.

## 5. THE NAHM TRANSFORM AND NAHM'S EQUATIONS

An alternative point of view on monopoles comes via Nahm's adaption of the Atiyah, Drinfeld, Hitchin, Manin construction of instantons [20]. Nahm considers a Dirac operator on  $\mathbb{R}^3$  coupled to the monopole. In more detail let  $\sigma^i$  be an orthonormal basis for the Lie algebra  $\mathfrak{su}(2)$ . This particular  $\mathfrak{su}(2)$  should be regarded as different to the monopole  $\mathfrak{su}(2)$  in which the connection and Higgs field take values. It is, in fact, the Lie algebra of the spin group of the group of rotations of  $\mathbb{R}^3$ . The Dirac operator  $D_z$  is defined by

$$D_z = \sum_{i=1}^3 \sigma^i \nabla_i - (\Phi + iz)$$

and acts on  $\mathbb{C}^2 \otimes \mathbb{C}^2$  valued functions on  $\mathbb{R}^3$ . The first  $\mathbb{C}^2$  is the spin space on which the  $\sigma^i$  act and the second is the space on which the  $A_i$  and  $\Phi$  act. Here  $z$  is any real number. We also define an adjoint

$$D_z^* = \sum_{i=1}^3 \sigma^i \nabla_i + (\Phi + iz).$$

If we compute the composition the Bogomolny equations show us that

$$D_z D_z^* = \sum_{i=1}^3 \nabla_i \nabla_i - (\Phi + iz)(\Phi + iz)$$

which is a positive operator and hence has no  $L^2$  kernel. From this we conclude that  $D_z^*$  has no  $L^2$  kernel. An  $L^2$  index theorem of Callias shows that  $D_z$  has index  $k$  if  $-1 < z < 1$  and 0 otherwise. Hence it follows that  $D_z$  has a  $k$  dimensional  $L^2$  kernel  $N_z$  if  $-1 < z < 1$ . The point of view we wish to adopt is that  $N_z$  is a  $k$  dimensional vector bundle over the interval  $(-1, 1)$ .



We are interested in sections of this vector bundle, that is functions

$$\psi: (-1, 1) \times \mathbb{R}^3 \rightarrow \mathbb{C}^2 \otimes \mathbb{C}^2.$$

which satisfy  $D_z \psi(z, x) = 0$  for every  $z \in (-1, 1)$ . Choose  $k$  of these  $\psi^1, \dots, \psi^k$  so that for each  $z$  they span  $N_z$ . Moreover choose them so that they are orthonormal

$$\int_{\mathbb{R}^3} (\psi^i, \psi^j) d^3x = \delta^{ij}$$

and satisfy

$$\int_{\mathbb{R}^3} \left( \psi^i, \frac{\partial \psi^j}{\partial z} \right) d^3x = 0$$

for all  $i, j = 1 \dots, k$ . Here, of course,  $\delta^{ij}$  is the Kronecker delta which is zero unless  $i = j$  when it is one. Notice that there is no obstruction to satisfying these extra conditions. We can use Gramm-Schmidt orthogonalisation for the first and the second is just solving an ordinary differential equation on  $(-1, 1)$ .

Now we define three  $k$  by  $k$  matrix functions of  $z$  by

$$T_a^{ij}(z) = \int_{\mathbb{R}^3} (\psi^i, x^a \psi^j) d^3x$$

for  $i, j = 1 \dots, k$  and  $a = 1, 2, 3$ . The remarkable thing about this Nahm transformation is that these matrix valued functions satisfy some simple ordinary differential equations, called Nahm's equations

$$\begin{aligned} \frac{dT_1}{dz} &= [T_2, T_3] \\ \frac{dT_2}{dz} &= [T_3, T_1] \\ \frac{dT_3}{dz} &= [T_1, T_2]. \end{aligned}$$

It is possible to cast Nahm's equations into Lax form and solve them by the Krichever method. We will consider how to do this as it connects us again with spectral curves.

Define

$$A(\zeta) = (T_1 + iT_2) + 2T_3\zeta + (-T_1 + iT_2)\zeta^2$$

and  $A_+(\zeta) = iT_3 - (iT_1 + T_2)\zeta$ . Then Nahm's equations are equivalent to

$$\frac{dA}{dz} = [A_+, A]$$

which is in Lax form. Now consider the curve  $S_z$  in  $\mathbb{C} \times \mathbb{C}$  defined by

$$(12) \quad \det(\eta - A(\zeta)) = 0.$$

We have

**Proposition 5.1.** *The curve  $S_z$  is independent of  $z$ .*

*Proof.* Choose an eigenvector  $v$  of  $A(\zeta)$  with eigenvalue  $\lambda$  at some  $z = z_0$ . Now evolve  $v$  with the differential equation

$$\frac{dv}{dz} = A_+ v$$

and consider

$$\begin{aligned}\frac{d(Av)}{dz} &= A_+Av - AA_+v + AA_+v \\ &= A_+(Av).\end{aligned}$$

It follows that

$$\frac{d}{dz}(Av - \lambda v) = 0$$

and because  $Av - \lambda v$  vanishes at  $z_0$  it must vanish everywhere. We conclude that the eigenvalues of  $A(\zeta)$  are independent of  $z$  and hence that  $S_z$  is independent of  $z$ .  $\square$

If we identify the  $(\eta, \zeta)$  in (12) with the co-ordinates (9) on mini-twistor space we realise  $S_z$  as a curve in mini-twistor space. This is, in fact, a natural thing to do. It is a remarkable fact [9] that the curve  $S = S_z$  defined via Nahm's transform is the same as the spectral curve of the monopole defined in Section 3.

Standard methods from integrable systems can be used to solve Nahm's equations using the curve  $S$  and some additional structure. Indeed in [7] Hitchin uses this approach to determine exactly which spectral curves correspond to monopoles.

One of the important properties of Nahm's equations is that it is straightforward to define a monopole from a solution of Nahm's equations plus some boundary conditions. Given such a solution and a point  $(x^1, x^2, x^3)$  in  $\mathbb{R}^3$  we define

$$T = \sum_{i=1}^3 T_i \sigma^i$$

and

$$x = \sum_{i=1}^3 x^i \sigma^i.$$

Now define  $E_x$  to be the  $L^2$  kernel of the operator

$$D_x = \frac{d}{dz} - T - x.$$

It can be shown that this is two dimensional. We define the connection and Higgs field by choosing an orthonormal basis  $(v_1, v_2)$  for  $E_x$  and letting

$$\Phi(v_a) = \sum_{b=1}^2 \int_{-1}^1 (v_b, zv_a) dz$$

and

$$A_i(v_a) = \sum_{b=1}^2 \int_{-1}^1 \left( v_b, \frac{dv_a}{dx^i} \right) dz.$$

This  $(A, \Phi)$  define a monopole if the  $T_i$  satisfy Nahm's equations with the appropriate boundary conditions. Moreover if the solution of Nahm's equations came from a monopole this construction returns the same monopole. For more details see [20].

## 6. RATIONAL MAPS

Although spectral curves are a useful invariant of monopoles it is very difficult given a spectral curve, for example a polynomial as in (11), to determine if it is the spectral curve of a monopole. Another approach to monopoles is the *rational map* of a monopole. By a rational map we mean a meromorphic function  $R(z): \mathbb{C} \rightarrow \mathbb{C}$  of the form

$$R(z) = \frac{p(z)}{q(z)}$$

where  $p$  and  $q$  are polynomials with no common divisor. We will be interested first in *based* rational maps, that is those which send  $\infty$  to 0 and have degree  $k$ . For a degree  $k$  based rational map we require that  $q$  is monic of degree  $k$  and that  $p$  has degree strictly less than  $k$ . Let  $\mathcal{R}_k^b$  be the set of all such maps then we have

**Theorem 6.1** (Donaldson [4]). *If we choose an orthogonal splitting of  $\mathbb{R}^3$  as  $\mathbb{C} \times \mathbb{R}$  then there is a diffeomorphism*

$$\mathcal{M}_k \simeq \mathcal{R}_k^b.$$

We shall indicate in a moment how to construct a rational map from a monopole. It is important to note that the inverse construction from rational map to monopole is not known. All that is known is that there is a map  $\mathcal{M}_k \rightarrow \mathcal{R}_k^b$  which can be shown to be a diffeomorphism. This is to be contrasted with the case of spectral curves where, although we have not discussed it, an inverse construction is known and with Nahm's construction where the inverse construction is quite straightforward. It seems that what we gain by knowing that every rational map is the rational map of some monopole we lose by not knowing how to get back to a monopole given a rational map.

To define the rational map we choose first the orthogonal splitting. This amounts to choosing a direction in  $\mathbb{R}^3$  and an orthogonal plane. Now consider Hitchin's equation along each line parallel to that direction. Because of the framing we have

$$\lim_{t \rightarrow \infty} \Phi(0, 0, t) = \frac{1}{\sqrt{2}} \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}.$$

We choose a gauge so that

$$\lim_{t \rightarrow \infty} \Phi(x, y, t) = \frac{1}{\sqrt{2}} \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}$$

for every  $(x, y)$ . For each  $z = x + iy$  consider the one-dimensional subspace of  $\mathbb{C}^2$  of solutions decaying along the line  $(x, y, t)$  as  $t \rightarrow \infty$ . Up to scale this line is spanned by a vector  $(a(z), b(z))$ . It turns out that  $b(z)$  is a polynomial of degree  $k$  with roots  $\beta^1, \dots, \beta^k$ . We choose the unique polynomial  $p(z)$  of degree  $k-1$  such that  $p(\beta^i) = a(\beta^i)$  for all  $i = 1, \dots, k$ . Finally we define  $R(z) = p(z)/b(z)$ . Notice that as  $|x|^2 + |y|^2 \rightarrow \infty$  the solution which decays as  $t \rightarrow -\infty$  blows up as  $t \rightarrow \infty$  and hence  $a(z) \rightarrow 0$  so that map  $R(z)$  is based. Note also that the spectral lines of the form  $(x, y, t)$  have  $b(x + iy) = 0$  so that they correspond to the poles of the rational map.

While an extremely useful invariant of the monopole as every based rational corresponds to exactly one monopole there are two significant problems. One we have already noted: the inverse map from rational maps to monopoles is not known. Another is that the construction breaks symmetry by writing  $\mathbb{R}^3$  as  $\mathbb{C} \times \mathbb{R}$  and hence

the construction does not preserve the full group of rotations of  $\mathbb{R}^3$  but instead the group of rotations around the axis which determines the splitting  $\mathbb{C} \times \mathbb{R}$ .

There is an alternative approach to rational maps due to Jarvis [13, 14] which avoids the problem of symmetry breaking. The rational map is also simpler to define. For this we pick a point in  $\mathbb{R}^3$  and consider Hitchin's equation along each (oriented) line through that point. Parametrise the lines through the point by  $\zeta$  and the distance along the line by  $t$  then the solution that decays in the forward direction is given by

$$s(\zeta, t) = \begin{pmatrix} s^0(\zeta, t) \\ s^1(\zeta, t) \end{pmatrix}.$$

The Jarvis rational map is then defined by

$$R(\zeta) = \frac{s^0(\zeta, 0)}{s^1(\zeta, 0)}.$$

To obtain a gauge invariant correspondence we note that the rational map is only determined up to the action of  $SU(2)$  by fractional linear transformations. Then Jarvis proves:

**Theorem 6.2** (Jarvis [13, 14]). *If we choose a point in  $\mathbb{R}^3$  there is a diffeomorphism*

$$\mathcal{M}_k/S^1 \simeq \mathcal{R}_k/SU(2).$$

Here  $\mathcal{R}_k$  is the set of all rational maps of degree  $k$  not just based and the space  $\mathcal{M}_k/S^1$  is the unframed moduli space, that is the moduli space divided by the action of the circle acting by constant gauge transformations.

The precise relationship between the Jarvis rational map and the Donaldson rational map has not yet been determined but it is believed that by moving the point determining the Jarvis map out to infinity along a line, and suitably rescaling the map, that the Donaldson map will be recovered.

## 7. CONCLUSION

We conclude by considering three significant omissions and generalisations.

Firstly the moduli space has a natural hyperkaehler metric. The physical interpretation of this is as follows. Consider the configuration space  $\mathcal{C}$  of all pairs  $(A, \Phi)$  satisfying the boundary conditions of a monopole. Inside  $\mathcal{C}$  we have the space  $\mathcal{C}_0$  of solutions of the Bogomolny equations and the monopole moduli space is the quotient of  $\mathcal{C}_0$  by the group of all gauge transformations. A solution of the full Yang-Mills-Higgs equations describes a curve  $(A(t), \Phi(t))$  in  $\mathcal{C}$  where the  $t$  parameter is time. Monopoles, points on  $\mathcal{C}_0$ , are static solutions of the full Yang-Mills-Higgs equations and the minima of a potential energy functional on  $\mathcal{C}$ . If we start with a point  $(A, \Phi)$  on  $\mathcal{C}_0$  and give it a small push then, by conservation of energy, it should describe a path staying close to  $\mathcal{C}_0$  and solving the full Yang-Mills-Higgs equations. In the limit, as the size of the push becomes zero, these paths determine geodesics of the natural hyperkaehler metric.

It follows that the geodesics on the monopole moduli space of this metric should approximate the motion of slowly moving monopoles. If we start with a monopole near the edge of the moduli space, which can be interpreted as a collection of  $k$  particles, and follow it along a geodesic until it emerges again into the region where it can be regarded as  $k$  particles we will have described a scattering process which should be an approximation to true  $k$ -monopole scattering. This idea is due to

Manton [16] and in [2] Atiyah and Hitchin describe the metric on the moduli space of charge two monopoles and certain of its geodesics and interpret the results in terms of two particle monopole scattering.

Secondly it is possible to generalise all the present discussion to other compact, simple Lie groups although the Nahm correspondence only works simply for the classical Lie groups. See for example, [17, 10, 11]. It is also possible to generalise to the loop group and loop group monopoles or calorons can be interpreted as instantons on  $S^1 \times \mathbb{R}^3$  [5].

Thirdly the underlying space  $\mathbb{R}^3$  can be replaced by hyperbolic three space and many of the results here carried over. This idea is due to Atiyah [1] and further developed in [3, 18] and [19].

## REFERENCES

- [1] M.F. Atiyah, *Magnetic Monopoles in hyperbolic spaces*, in Proceedings of Bombay Colloquium 1984 on vector bundles in algebraic varieties (Oxford University Press), pp. 1–34, 1987.
- [2] M.F. Atiyah and N.J. Hitchin, *The geometry and dynamics of magnetic monopoles*, Princeton Univ. Press, Princeton, NJ, 1988.
- [3] P.J. Braam and D.M. Austin, *Boundary values of hyperbolic monopoles*, Nonlinearity **3** (1990), no. 3, 809–823.
- [4] S.K. Donaldson, *Nahm’s equations and the classification of monopoles*, Commun. Math. Phys. **96** (1984), 387–407.
- [5] H. Garland and M.K. Murray, *Kac-Moody monopoles and periodic instantons*, Commun. Math. Phys. **120** (1988), 335–351.
- [6] N.J. Hitchin, *On the construction of monopoles*, Commun. Math. Phys. **89** (1983), 145–190.
- [7] N.J. Hitchin, *Monopoles and Geodesics*, Commun. Math. Phys. **83** (1982), 579–602.
- [8] N.J. Hitchin, N.S. Manton and M.K. Murray, *Symmetric Monopoles*, Nonlinearity **8** (1995), 661–692.
- [9] N.J. Hitchin and M.K. Murray, *Spectral curves and the ADHM method*, Commun. Math. Phys. **114** (1988), 463–474.
- [10] J. Hurtubise and M.K. Murray, *On the construction of monopoles for the classical groups*, Commun. Math. Phys. **122** (1989), 35–89.
- [11] J.C. Hurtubise and M.K. Murray, *Monopoles and their spectral data*, Commun. Math. Phys. **133** (1990), 487–508.
- [12] A. Jaffe and C. Taubes, *Vortices and monopoles*, Birkhäuser, Boston, MA, 1980.
- [13] S. Jarvis, *Construction of Euclidean monopoles*, Proc. London Math. Soc. (3) **77** (1998), 193–214.
- [14] S. Jarvis, *Euclidean monopoles and rational maps*, Proc. London Math. Soc. (3) **77** (1998), 170–192.
- [15] N.S. Manton, *The force between ‘t Hooft-Polyakov monopoles*, Nucl. Phys. **B126** (1977), 525–541.
- [16] N.S. Manton, *A remark on the scattering of BPS monopoles*, Phys. Lett. **110B** (1982), 54–56.
- [17] M.K. Murray, *Non-abelian Magnetic Monopoles*, Commun. Math. Phys. **96** (1984), 539–565.
- [18] M.K. Murray and M.A. Singer, *Spectral curves of non-integral hyperbolic monopoles*, Nonlinearity **9** (1996), 973–997.
- [19] M.K. Murray and M.A. Singer, *On the complete integrability of the discrete Nahm equations* Communications in Mathematical Physics, **210**, no. 2, (2000) 497–519.
- [20] W. Nahm, *The construction of all self-dual monopoles by the ADHM method*, in Monopoles in Quantum Field Theory, Proceedings of the monopole meeting in Trieste 1981, World Scientific, Singapore (1982).
- [21] M.K. Prasad and C.M. Sommerfield, *Exact classical solutions for the ‘t Hooft monopole and Julia-Zee dyon*, Phys. Rev. Lett. **35** (1975), 760–762.
- [22] P. Sutcliffe, *BPS monopoles*, Int. J. Mod. Phys. **A12** (1997), 4663–4706.
- [23] E. Weinberg, *Parameter counting for multimonopole solutions*, Phys. Rev. **D20** (1979), 936–944.

DEPARTMENT OF PURE MATHEMATICS, UNIVERSITY OF ADELAIDE, ADELAIDE, SA 5005, AUSTRALIA

*E-mail address:* `mmurray@maths.adelaide.edu.au`